

UNCLASSIFIED

AD 261 343

*Reproduced
by the*

**ARMED SERVICES TECHNICAL INFORMATION AGENCY
ARLINGTON HALL STATION
ARLINGTON 12, VIRGINIA**



UNCLASSIFIED

DISCLAIMER NOTICE

**THIS DOCUMENT IS BEST QUALITY
PRACTICABLE. THE COPY FURNISHED
TO DTIC CONTAINED A SIGNIFICANT
NUMBER OF PAGES WHICH DO NOT
REPRODUCE LEGIBLY.**

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

CATALOGUE No. 11A
AS AD No. 261343

TECHNICAL REPORT FRL-TR-38

EPR OBSERVATION OF NH_4^+ FORMED BY X-RAY
IRRADIATION OF AMMONIUM PERCHLORATE
CRYSTALS


JAMES S. HYDE, INSTRUMENT DIVISION,
VARIAN ASSOCIATES, PALO ALTO, CALIFORNIA

ELI S. FREEMAN, PICATINNY ARSENAL

AUGUST 1961



FELTMAN RESEARCH LABORATORIES
PICATINNY ARSENAL
DOVER, N. J.



OMS 5530.11.558

DEPT. OF THE ARMY PROJECT 5504-01-027

COPY 22

2161.4-2
XEROX

DISPOSITION

Destroy this report when it is no longer needed.
Do not return.

ASTIA AVAILABILITY NOTICE

Qualified requesters may obtain copies of this report
from ASTIA.

EPR OBSERVATION OF NH_3^+ FORMED BY X-RAY IRRADIATION
OF AMMONIUM PERCHLORATE CRYSTALS

by

James S. Hyde,
Instrument Division, Varian Associates, Palo Alto, California
and
Eli S. Freeman, Picatinny Arsenal

August 1961

Feltman Research Laboratories
Picatinny Arsenal
Dover, N. J.

Technical Report FRL-TR-38

OMS 5530.11.558

Dept of the Army Project 5504-01-027

Approved:

Sage
S. SAGE
Chief, Pyrotechnics
Laboratory

TABLE OF CONTENTS

	Page
Abstract	1
Introduction	2
Sample Preparation	2
Electron Paramagnetic Resonance Apparatus	2
Results and Discussion	2
References	4
Distribution List	7
Figures	
1	Room temperature EPR spectrum from a single crystal of irradiated ammonium perchlorate, and the reconstructed spectrum from NH_3^+ assuming isotropic hyperfine interaction
5	
2	EPR spectra at $+100^\circ\text{C}$, $+25^\circ\text{C}$, -100°C , and -180°C from a crystalline powder of irradiated ammonium perchlorate
6	

7

ABSTRACT

An investigation of irradiated ammonium perchlorate was conducted by electron spin resonance. A new specie, the NH_3^+ radical was discovered. This ion radical appears to be stable up to at least 125°C.

INTRODUCTION

Profound changes in the decomposition characteristics or chemical reactivity of ammonium perchlorate crystals following X-ray and gamma ray irradiation have been reported (Refs 1 and 2). In an effort to understand the nature of the radiation damage in these crystals, we have undertaken an investigation using electron paramagnetic resonance (EPR) techniques. One would expect, with these techniques, to observe those damage sites in the crystal which happen to contain unpaired electron spins.

SAMPLE PREPARATION

The ammonium perchlorate crystals used in this work were doubly recrystallized from triply distilled water. The original material was certified reagent grade salt obtained from the Fisher Scientific Company. Both crystalline powders and larger crystals of several cubic millimeters were prepared. The samples were irradiated with an exposure dose of 10^6 roentgens at a rate of 2×10^5 roentgens per hour by means of an OEG-50 X-ray tube with a molybdenum target.

ELECTRON PARAMAGNETIC RESONANCE APPARATUS

The electron paramagnetic resonance experiments were performed using the Varian V4501 EPR spectrometer. This instrument employs 100 kc field modulation and a microwave frequency of 9.5 kmc. A Varian V4547 variable temperature accessory permitted studies over a continuous range of temperature.

RESULTS AND DISCUSSION

Figure 1 shows the EPR spectrum obtained at room temperature from a single crystal of irradiated ammonium perchlorate. The spectrum is what might be expected from hyperfine interaction with one nucleus of spin 1 and three equivalent nuclei of spin $1/2$, and almost certainly arises from an unpaired electron spin strongly localized on an NH_3 molecule. Contact with the nitrogen nucleus splits the spectrum into three lines of equal spacing and intensity, and contact with the three equivalent

protons splits each of these three lines into a quartet of equally spaced lines with intensity ratios of 1:3:3:1. This analysis is illustrated in Figure 1. Presumably the radical is a positive ion, although the EPR technique would not distinguish it from a negative ion. No resonances other than that associated with the NH_3^+ radical were found in these crystals.

Several interesting observations can be made concerning the anisotropy and temperature dependence. Figure 2 shows spectra obtained with a powder sample at $+100^\circ\text{C}$, $+25^\circ\text{C}$, -100°C , and -180°C . At $+25^\circ\text{C}$, the spectra from the powder and the single crystal differ markedly, but at $+100^\circ\text{C}$ the spectra are nearly the same. We conclude that, at the high temperature, the NH_3^+ molecules are rotating rapidly and the anisotropy which causes the difference in spectra found at lower temperatures is being averaged out. The critical rotation frequency is $\gamma \overline{\Delta H}$, where γ is the gyromagnetic ratio of the electron dipole and $\overline{\Delta H}$ is an average shift in the position of the hyperfine lines because of the anisotropy. The separation of the lines at $+100^\circ\text{C}$ is therefore a measure of the isotropic hyperfine splitting. Contact with the nitrogen nucleus splits the lines by 18.1 oersteds and contact with the hydrogen nuclei splits the lines by 25.0 oersteds. The splittings were measured to an accuracy of one per cent by comparing them with the resonance of nitrosodisulphonate (peroxylamine disulphonate) ion $(\text{ON}(\text{SO}_3)_2)^\pm$, which is split into three lines 13.0 oersteds apart.

At -100°C (Fig 2), the central quartet corresponding to $M = 0$ for the nitrogen nucleus is well defined and of much greater intensity than the quartets corresponding to $M = \pm 1$. It is quite possible that the observed anisotropy arises from interaction with the quadrupole moment of the nitrogen nucleus, since the nitrogen splittings are so much more drastically affected than are the hydrogen splittings. Hydrogen, of course, has no quadrupole moment. It is also possible that the electron dipole has stronger anisotropic interaction with the dipole moment of the nitrogen nucleus than with the proton dipoles. In any event, the molecule is rotating slowly compared with $(\gamma \overline{\Delta H_N})^{-1}$, where $\overline{\Delta H_N}$ refers to the average anisotropic shift arising from interaction with the nitrogen nucleus. At still lower temperatures (-180°C), the spectrum from the

powder becomes asymmetric with respect to the center of the spectrum (Fig 2), although the single crystal spectrum remains symmetric. Anisotropy arising from interaction between the electron dipole and the proton dipoles enters the picture, and the molecule rotates slowly compared with $(\gamma \Delta H_P)^{-1}$ where the subscript "P" refers to the protons. Rogers and Pake³ have reported similar anisotropic effects when the vanadyl ion, VO^{++} , is in aqueous solution. The behavior of the anisotropy of the NH_3^+ resonance as a function of temperature suggests that freedom of the ion to rotate is frozen out in several fairly well defined steps.

The ammonia ion is quite stable with respect to time and temperature. No decay of the resonance intensity was noticed upon heating to $+125^\circ C$ or upon storing for several months at room temperature.

The resonance cannot be saturated at room temperature with the available microwave power (200 mw at the sample cavity); at lower temperatures, however, it is necessary to reduce the power level to avoid saturation. The g value of the center of gravity of the spectrum was measured by comparison with crystalline dpph ($g = 2.0036$) and found to be 2.0034 ± 0.0001 .

REFERENCES

1. E. S. Freeman and D. A. Anderson, J. Phys. Chem. **63**, 1344 (1959)
2. E. S. Freeman, D. A. Anderson, and J. J. Campisi, J. Phys. Chem. **64**, 1727 (1960)
3. R. N. Rogers and G. E. Pake, J. Chem. Phys. **33**, 1107 (1960)





Fig.1 Room temperature EPR spectrum from a single crystal of irradiated ammonium perchlorate, and the reconstructed spectrum from NH_3^+ assuming isotropic hyperfine interaction




Fig 2 EPR spectra at $+100^{\circ}\text{C}$, $+25^{\circ}\text{C}$, -100°C , and -180°C from a crystalline powder of irradiated ammonium perchlorate

DISTRIBUTION LIST

Copy No.

Commanding Officer Picatinny Arsenal ATTN: Technical Information Section Dover, N. J.	1 - 5
Chief of Ordnance Dept of the Army ATTN: ORDIM	6
ORDTS	7
ORDTB	8
Washington 25, D. C.	
Commanding General OSWAC Picatinny Arsenal ATTN: ORDSW-A	9
ORDSW-W	10
Dover, N. J.	
Commanding General Frankford Arsenal ATTN: Pitman-Dunn Laboratory	11
Bridge and Tacony Streets Philadelphia 37, Pennsylvania	
Commanding General Aberdeen Proving Ground ATTN: Dev and Proof Services	12
BRL	13
Maryland	
Armed Services Technical Information Agency Arlington Hall Station Arlington 12, Virginia	14 - 23
Bureau of Naval Weapons Dept of the Navy ATTN: Re2A	24 - 25
Washington 25, D. C.	

Copy No.

Commander
Naval Ordnance Laboratory
White Oak, Silver Spring
Maryland

26

Commander
Wright Air Development Division
ATTN: WCLFE-E
Wright-Patterson Air Force Base
Ohio

27

Commanding Officer
Redstone Arsenal
Alabama

28 - 29

Commander
Air Research and Development Command
Andrews Air Force Base
Washington 25, D. C.

30

Commanding Officer
Diamond Ordnance Fuze Laboratories
ATTN: Technical Reference Section
Connecticut Ave and Van Ness St, NW
Washington 25, D. C.

31

Commander
U. S. Naval Ordnance Test Station
China Lake, Inyokern
California

32 - 33

Army Research Office (Durham)
Box CM, Duke Station
Durham, North Carolina

34

Commanding General
Ordnance Ammunition Command
ATTN: ORDLY-R
Joliet, Illinois

35

Copy No.

Commanding Officer Longhorn Ordnance Works Marshall, Texas	36
Commanding Officer Jefferson Proving Ground Madison, Indiana	37
Director of Research and Development Dept of the Air Force Headquarters, USAF, DCS/D ATTN: AFDRD-EQ-3	38
Coordinating Member	39
ASCC Working Party 23 Washington 25, D. C.	40
Commanding General White Sands Missile Range New Mexico	41
Headquarters Continental Army Command Ft. Monroe, Virginia	42
Commanding Officer U. S. Army Ordnance Test Activity Yuma, Arizona	43
Commanding General Air Proving Ground Center Eglin Air Force Base Florida	44
Commanding Officer Naval Missile Center Point Mugu, California	45
National Aeronautics and Space Administration 1512 H St., NW Washington 25, D. C.	46

Copy No.

Chief, U. S. Army Research Office
Research Analysis Division
ATTN: Dr. Hoyt Lemons
Arlington Hall Station, Virginia

47

Assistant Secretary of Defense
(Research and Engineering)
Technical Library
Pentagon 3E1065
Washington 25, D. C.

48

Commander
Air Force Cambridge Research Center
ATTN: CROOTR, L. G. Hanscom Field
Bedford, Massachusetts

49

AD _____ Accession No. _____

Feltman Research Laboratories

Picatinny Arsenal, Dover, N. J.

EPR OBSERVATION OF NH_4^+ FORMED BY X-RAY IRRADIATION OF AMMONIUM PERCHLORATE CRYSTALS

James S. Hyde, Instrument Division, Varian Associates, Palo Alto, California and Eli S. Freeman, Picatinny Arsenal

Technical Report FRL-TR-38, August 1961, 10 pp, figures. DA Proj 5504-01-027, OMS 5530.11.558. Unclassified Report

An investigation of irradiated ammonium perchlorate was conducted by electron spin resonance. A new species, the NH_4^+ radical was discovered. This ion radical appears to be stable up to at least 100°C.

(over)

UNCLASSIFIED

1. Paramagnetic crystals—Spectrographic analysis

2. Ammonium perchlorate crystals—Effects of radiation

3. Ammonia ions

I. Hyde, James S.

II. Freeman, Eli S.

III. Title

IV. OMS 5530.11.558

V. DA Proj 5504-01-027

UNITERMS

Ammonium perchlorate

X-rays

UNCLASSIFIED

AD _____ Accession No. _____

Feltman Research Laboratories

Picatinny Arsenal, Dover, N. J.

EPR OBSERVATION OF NH_4^+ FORMED BY X-RAY IRRADIATION OF AMMONIUM PERCHLORATE CRYSTALS

James S. Hyde, Instrument Division, Varian Associates, Palo Alto, California and Eli S. Freeman, Picatinny Arsenal

Technical Report FRL-TR-38, August 1961, 10 pp, figures. DA Proj 5504-01-027, OMS 5530.11.558. Unclassified Report

An investigation of irradiated ammonium perchlorate was conducted by electron spin resonance. A new species, the NH_4^+ radical was discovered. This ion radical appears to be stable up to at least 100°C.

AD _____ Accession No. _____

Feltman Research Laboratories

Picatinny Arsenal, Dover, N. J.

EPR OBSERVATION OF NH_4^+ FORMED BY X-RAY IRRADIATION OF AMMONIUM PERCHLORATE CRYSTALS

James S. Hyde, Instrument Division, Varian Associates, Palo Alto, California and Eli S. Freeman, Picatinny Arsenal

Technical Report FRL-TR-38, August 1961, 10 pp figures. DA Proj 5504-01-027, OMS 5530.11.558. Unclassified Report

An investigation of irradiated ammonium perchlorate was conducted by electron spin resonance. A new species, the NH_4^+ radical was discovered. This ion radical appears to be stable up to at least 100°C.

(over)

UNCLASSIFIED

1. Paramagnetic crystals—Spectrographic analysis

2. Ammonium perchlorate crystals—Effects of radiation

3. Ammonia ions

I. Hyde, James S.

II. Freeman, Eli S.

III. Title

IV. OMS 5530.11.558

V. DA Proj 5504-01-027

UNITERMS

Ammonium perchlorate

X-rays

UNCLASSIFIED

UNCLASSIFIED

1. Paramagnetic crystals—Spectrographic analysis

2. Ammonium perchlorate crystals—Effects of radiation

3. Ammonia ions

I. Hyde, James S.

II. Freeman, Eli S.

III. Title

IV. OMS 5530.11.558

V. DA Proj 5504-01-027

UNITERMS

Ammonium perchlorate

X-rays

UNCLASSIFIED

UNCLASSIFIED
UNITERMS

Effect
Resonance methods
Ammonia
Ion
Hyde, J. S.
Freeman, E. S.
OMS 5530.11.558
DA Proj 5504-01-027

UNCLASSIFIED

UNCLASSIFIED
UNITERMS

Effect
Resonance methods
Ammonia
Ion
Hyde, J. S.
Freeman, E. S.
OMS 5530.11.558
DA Proj 5504-01-027

UNCLASSIFIED

UNCLASSIFIED
UNITERMS

Effect
Resonance methods
Ammonia
Ion
Hyde, J. S.
Freeman, E. S.
OMS 5530.11.558
DA Proj 5504-01-027

UNCLASSIFIED

UNCLASSIFIED
UNITERMS

Effect
Resonance methods
Ammonia
Ion
Hyde, J. S.
Freeman, E. S.
OMS 5530.11.558
DA Proj 5504-01-027

UNCLASSIFIED

AD _____ Accession No. _____
Feltman Research Laboratories
Picatinny Arsenal, Dover, N.J.
**EPR OBSERVATION OF NH_4^+ FORMED BY X-RAY
IRRADIATION OF AMMONIUM PERCHLORATE
CRYSTALS**
James S. Hyde, Instrument Division, Varian Associates,
Palo Alto, California and Eli S. Freeman, Picatinny
Arsenal

Technical Report FRL-TR-38, August 1961, 10 pp,
figures. DA Proj 5504-01-027, OMS 5530.11.558.
Unclassified Report

An investigation of irradiated ammonium perchlorate
was conducted by electron spin resonance. A new
species, the NH_3^+ radical was discovered. This ion
radical appears to be stable up to at least 100°C.

UNCLASSIFIED

1. Paramagnetic crystals--
Spectrographic
analysis
2. Ammonium perchlorate
crystals--Effects of
radiation
3. Ammonia ions
- I. Hyde, James S.
- II. Freeman, Eli S.
- III. Title
- IV. OMS 5530.11.558
- V. DA Proj 5504-01-
027
- UNITERMS
Ammonium perchlorate
X-rays

(over)

AD _____ Accession No. _____
Feltman Research Laboratories
Picatinny Arsenal, Dover, N.J.
**EPR OBSERVATION OF NH_4^+ FORMED BY X-RAY
IRRADIATION OF AMMONIUM PERCHLORATE
CRYSTALS**
James S. Hyde, Instrument Division, Varian Associates,
Palo Alto, California and Eli S. Freeman, Picatinny
Arsenal

Technical Report FRL-TR-38, August 1961, 10 pp,
figures. DA Proj 5504-01-027, OMS 5530.11.558.
Unclassified Report

An investigation of irradiated ammonium perchlorate
was conducted by electron spin resonance. A new
species, the NH_3^+ radical was discovered. This ion
radical appears to be stable up to at least 100°C.

UNCLASSIFIED

1. Paramagnetic crystals--
Spectrographic
analysis
2. Ammonium perchlorate
crystals--Effects of
radiation
3. Ammonia ions
- I. Hyde, James S.
- II. Freeman, Eli S.
- III. Title
- IV. OMS 5530.11.558
- V. DA Proj 5504-01-
027
- UNITERMS
Ammonium perchlorate
X-rays

UNCLASSIFIED

AD _____ Accession No. _____
Feltman Research Laboratories
Picatinny Arsenal, Dover, N.J.
**EPR OBSERVATION OF NH_4^+ FORMED BY X-RAY
IRRADIATION OF AMMONIUM PERCHLORATE
CRYSTALS**
James S. Hyde, Instrument Division, Varian Associates,
Palo Alto, California and Eli S. Freeman, Picatinny
Arsenal

Technical Report FRL-TR-38, August 1961, 10 pp
figures. DA Proj 5504-01-027, OMS 5530.11.558.
Unclassified Report

An investigation of irradiated ammonium perchlorate
was conducted by electron spin resonance. A new
species, the NH_3^+ radical was discovered. This ion
radical appears to be stable up to at least 100°C.

(over)

AD _____ Accession No. _____
Feltman Research Laboratories
Picatinny Arsenal, Dover, N.J.
**EPR OBSERVATION OF NH_4^+ FORMED BY X-RAY
IRRADIATION OF AMMONIUM PERCHLORATE
CRYSTALS**
James S. Hyde, Instrument Division, Varian Associates,
Palo Alto, California and Eli S. Freeman, Picatinny
Arsenal

Technical Report FRL-TR-38, August 1961, 10 pp,
figures. DA Proj 5504-01-027, OMS 5530.11.558.
Unclassified Report

An investigation of irradiated ammonium perchlorate
was conducted by electron spin resonance. A new
species, the NH_3^+ radical was discovered. This ion
radical appears to be stable up to at least 100°C.

UNCLASSIFIED

1. Paramagnetic crystals--
Spectrographic
analysis
2. Ammonium perchlorate
crystals--Effects of
radiation
3. Ammonia ions
- I. Hyde, James S.
- II. Freeman, Eli S.
- III. Title
- IV. OMS 5530.11.558
- V. DA Proj 5504-01-
027
- UNITERMS
Ammonium perchlorate
X-rays

UNCLASSIFIED

1. Paramagnetic crystals--
Spectrographic
analysis
2. Ammonium perchlorate
crystals--Effects of
radiation
3. Ammonia ions
- I. Hyde, James S.
- II. Freeman, Eli S.
- III. Title
- IV. OMS 5530.11.558
- V. DA Proj 5504-01-
027
- UNITERMS
Ammonium perchlorate
X-rays

UNCLASSIFIED

UNCLASSIFIED
UNITERMS

Effect
Resonance methods
Ammonia
Ion
Hyde, J. S.
Freeman, E. S.
OMS 5530.11.558
DA Proj 5504-01-027

UNCLASSIFIED

UNCLASSIFIED
UNITERMS

Effect
Resonance methods
Ammonia
Ion
Hyde, J. S.
Freeman, E. S.
OMS 5530.11.558
DA Proj 5504-01-027

UNCLASSIFIED

UNCLASSIFIED
UNITERMS

Effect
Resonance methods
Ammonia
Ion
Hyde, J. S.
Freeman, E. S.
OMS 5530.11.558
DA Proj 5504-01-027

UNCLASSIFIED

UNCLASSIFIED
UNITERMS

Effect
Resonance methods
Ammonia
Ion
Hyde, J. S.
Freeman, E. S.
OMS 5530.11.558
DA Proj 5504-01-027

UNCLASSIFIED

UNCLASSIFIED

UNCLASSIFIED
UNITERMS

Effect
Resonance methods
Ammonia
Ion
Hyde, J. S.
Freeman, E. S.
OMS 5530.11.558
DA Proj 5504-01-027

UNCLASSIFIED

UNCLASSIFIED
UNITERMS

Effect
Resonance methods
Ammonia
Ion
Hyde, J. S.
Freeman, E. S.
OMS 5530.11.558
DA Proj 5504-01-027

UNCLASSIFIED

UNCLASSIFIED
UNITERMS

Effect
Resonance methods
Ammonia
Ion
Hyde, J. S.
Freeman, E. S.
OMS 5530.11.558
DA Proj 5504-01-027

UNCLASSIFIED

UNCLASSIFIED
UNITERMS

Effect
Resonance methods
Ammonia
Ion
Hyde, J. S.
Freeman, E. S.
OMS 5530.11.558
DA Proj 5504-01-027

UNCLASSIFIED